

WHAT TO EXPECT WITH YORVIPATH

YOUR GUIDE TO STARTING TREATMENT



Lynette, **actual patient**

IMPORTANT SAFETY INFORMATION AND USE

What is YORVIPATH® used for?

YORVIPATH is a prescription medicine used to treat adults with low parathyroid hormone (PTH) (hypoparathyroidism).

- It is not known if YORVIPATH is safe and effective in people who have been recently diagnosed with hypoparathyroidism after surgery, or if it is started in people with low levels of calcium in the blood.
- It is not known if YORVIPATH is safe and effective in children.
- YORVIPATH should not be used in children and young adults whose bones are still growing.

Who should not take YORVIPATH?

Do not use YORVIPATH if you have had a severe allergic reaction to palopegteriparatide or any of the other ingredients in YORVIPATH.

Please see Important Safety Information throughout. Please click to read or print the [Prescribing Information](#) and [Medication Guide](#) for YORVIPATH.

YORVIPATH DELIVERS CONTINUOUS PARATHYROID HORMONE (PTH) OVER 24 HOURS

YORVIPATH is a once-daily injection that is designed to help provide your body with the PTH it needs



YORVIPATH CAN HELP:



Maintain normal blood calcium levels



Maintain normal movement of calcium through your bones, blood, and vital organs



Activate vitamin D to help with calcium absorption

IMPORTANT SAFETY INFORMATION AND USE

What is the most important information I should know about YORVIPATH?

YORVIPATH may cause serious side effects, including:

- **Allergic (hypersensitivity) reactions, including anaphylaxis.** Stop taking YORVIPATH and get medical help right away if you have any of the following symptoms of an allergic reaction: swelling of your face, lips, mouth or tongue, breathing problems, fainting, dizziness, feeling lightheaded (low blood pressure), fast heartbeat, itching, rash or hives.
- **High levels of calcium in the blood (hypercalcemia).** YORVIPATH can cause some people to have higher blood calcium levels than normal. Your healthcare provider should check your blood calcium before you start and during your treatment with YORVIPATH. Tell your healthcare provider right away if you have nausea, vomiting, dizziness, feeling thirsty, confusion, muscle weakness, or irregular heartbeat. Hypercalcemia is more likely to occur within the first 3 months of starting YORVIPATH, but it may occur at any time.

Please see Important Safety Information throughout. Please click to read or print the [Prescribing Information](#) and [Medication Guide](#) for YORVIPATH.

Yorvipath®
paleopecteriparatide
Injection 300 mcg/mL

Starting with the basics

EACH PACKAGE IS SHIPPED FROM YOUR SPECIALTY PHARMACY AND COMES WITH:

- Two pre-filled pens
- Twenty-eight needles
- Two spare needles



TOOLS AND RESOURCES TO HELP YOU STAY ON TRACK



Enroll in the Ascendis Signature Access Program (A-S-A-P) for personalized support and a starter kit



Use the treatment journal in your kit to track injections

IMPORTANT SAFETY INFORMATION AND USE

What is the most important information I should know about YORVIPATH? (cont'd)

YORVIPATH may cause serious side effects, including:

- **Low levels of calcium in the blood (hypocalcemia).** People who stop using, miss, or change a dose of YORVIPATH may have an increased risk of low blood calcium levels, but hypocalcemia may occur at any time. Tell your healthcare provider right away if you have tingling in your fingertips, toes, lips or tongue, muscle spasms or cramps, oral numbness, depression, have problems thinking or remembering, abnormal heart rhythms, or seizures.

Please see Important Safety Information throughout. Please click to read or print the [Prescribing Information](#) and [Medication Guide](#) for YORVIPATH.

Yorvipath®
palopegteriparatide
Injection 300 mcg/mL

Getting started with your pen



PREPARATION

- You will be taught how to inject YORVIPATH by your doctor or nurse before your first dose
- Until first use, store your pens in the refrigerator between 36°F to 46°F (2°C to 8°C)



FIRST USE

- Take a new pen out of the refrigerator 20 minutes before use to allow the medicine to reach room temperature
- For up to 14 days after your first use, store your pen at room temperature, up to 86°F (30°C)



DISPOSAL

- Discard your pen after 14 days, even if it still has medicine inside it
- Check your state guidelines for instructions on how to properly dispose of your pen and needles



TRAVEL

- Traveling with YORVIPATH is simple when you follow the travel checklist



Download the travel checklist



For personalized support, call **1-844-442-7236** from 8 AM to 8 PM ET, Monday through Friday.

IMPORTANT SAFETY INFORMATION AND USE

What is the most important information I should know about YORVIPATH? (cont'd)


YORVIPATH may cause serious side effects, including:

- **Possible bone cancer (osteosarcoma).** Tell your healthcare provider right away if you have pain in any areas of your body that does not go away or any new or unusual lumps or swelling under your skin that is tender to touch. These are some of the signs and symptoms of osteosarcoma and your healthcare provider may need to do further tests.


Please see Important Safety Information throughout. Please click to read or print the [Prescribing Information](#) and [Medication Guide](#) for YORVIPATH.

Yorvipath®
palopegteriparatide
Injection 300 mcg/mL

Taking a closer look



*Depending on your pen, the dose window will show 6, 9, or 12 mcg; 15, 18, or 21 mcg; or 24, 27, or 30 mcg.



IMPORTANT SAFETY INFORMATION AND USE

Who should not take YORVIPATH?

Do not use YORVIPATH if you have had a severe allergic reaction to palopegteriparatide or any of the other ingredients in YORVIPATH.

Please see Important Safety Information throughout. Please click to read or print the [Prescribing Information](#) and [Medication Guide](#) for YORVIPATH.

Yorvipath®
palopegteriparatide
Injection 300 mcg/mL

Finding your individualized dose

Achieving your individualized YORVIPATH dose requires a process called titration. During this process, your dose is slowly adjusted based on your lab results. Your doctor will develop your titration schedule and work with you along the way.

Finding the YORVIPATH dose that works for you can take some time, so don't get discouraged.

THE GOAL OF THIS PROCESS



Find the dose needed to maintain blood calcium levels in the normal range without active vitamin D and calcium

Talk to your doctor if you have any questions throughout the process. For more information about individualized dosing, visit YORVIPATH.com.

Chris,
actual patient



IMPORTANT SAFETY INFORMATION AND USE

What should I tell my healthcare provider before taking YORVIPATH?

Before using YORVIPATH, tell your healthcare provider about all of your medical conditions, including if you:

- are at higher risk of a type of bone cancer called osteosarcoma. This is especially important:
 - if you have a bone disease that increases your risk of developing osteosarcoma (including if you have Paget's disease).
 - if a blood test shows that you have unexplained increases in bone alkaline phosphatase.
 - if you have cancer of the bones or other cancer that has spread to your bones.
 - if you are having or have had radiation therapy to the skeleton.

Please see Important Safety Information throughout. Please click to read or print the [Prescribing Information](#) and [Medication Guide](#) for YORVIPATH.

Yorvipath[®]
paleogepteriparatide
Injection 300 mcg/mL

The steps you'll follow



EVALUATE

Two weeks before you start YORVIPATH, your doctor will make sure your blood calcium and vitamin D levels are within the recommended range.



START

Your doctor will start you on YORVIPATH (the recommended starting dose is 18 micrograms, once daily).

- Since YORVIPATH is designed to help provide your body with the PTH it needs, your doctor will adjust your calcium and active vitamin D doses based on your lab results



ADJUST

Everyone progresses at a different rate during this time.

- Your YORVIPATH dose may be adjusted according to your lab results until the individualized dose for your body is found
- Your dose of active vitamin D and/or calcium may continue to be adjusted in this phase
- 7-10 days after each adjustment, your blood calcium levels will be checked



CONTINUE

Your individualized dose is the amount of YORVIPATH you will continue to take once your blood calcium levels are within a normal range.

- Your blood calcium levels will continue to be assessed every 4-6 weeks throughout your treatment with YORVIPATH, as your dose may still need to be adjusted over time
- You may still need to take some amount of calcium according to your doctor's recommendation

IMPORTANT SAFETY INFORMATION AND USE

What should I tell my healthcare provider before taking YORVIPATH? (cont'd)

Before using YORVIPATH, tell your healthcare provider about all of your medical conditions, including if you:

- are at higher risk of a type of bone cancer called osteosarcoma. This is especially important:
 - if you are affected with a condition that runs in your family that can increase your chance of getting cancer in your bones.

Please see Important Safety Information throughout. Please click to read or print the [Prescribing Information](#) and [Medication Guide](#) for YORVIPATH.

Yorvipath[®] 
palopegteriparatide
Injection 300 mcg/mL

Follow the P.A.T.H. to injection

Keep these 4 steps in mind when you inject YORVIPATH.*†

P

PREPARE

- Wash your hands
- Gather your pen, needle, and alcohol wipes
- Check the expiration date and ensure the medicine is clear
- Click the needle straight onto the pen, then screw it in until secure

A

ASSESS

- Choose and clean your injection site
- Align your dose by turning the selector clockwise until it matches your prescribed amount

T

TREAT

- Insert the needle into the skin at your injection site
- Push button down until the dose selector rotates back to the “•” symbol
- Hold steady for 5 seconds before removing

H

HANDLE & HOUSE

- Reattach the outer needle cover to carefully remove the needle
- Unscrew the needle and discard it according to local regulations
- Snap the cap back onto the pen to protect it from light

Once you start using a pen, it must be discarded after 14 days, even if there is medicine left inside.

*These points serve only as some reminders for YORVIPATH. You should follow direction from your doctor, fully review the Instructions for Use provided with YORVIPATH, and follow the directions provided during your injection training.

†Before starting YORVIPATH, be sure to contact your doctor. You may require adjustments to your current medication.

IMPORTANT SAFETY INFORMATION AND USE

What should I tell my healthcare provider before taking YORVIPATH? (cont'd)

Before using YORVIPATH, tell your healthcare provider about all of your medical conditions, including if you:

- take medicines that contain digoxin, are used to treat osteoporosis, or can affect calcium levels in your blood.

Please see Important Safety Information throughout. Please click to read or print the [Prescribing Information](#) and [Medication Guide](#) for YORVIPATH.

Yorvipath[®]
palopegteriparatide
Injection 300 mcg/mL

Important injection reminders

CHOOSING AN INJECTION SITE




Inject YORVIPATH into the belly (abdomen) or in the front of the thigh



Choose a new site for each injection



Avoid injecting where skin is red, swollen, or scarred



It is important to take YORVIPATH at the same time every day, whether you choose to inject during the day or at night.



Ready to get started injecting?

Review detailed instructions and watch the step-by-step video at [YORVIPATH.com](https://www.yorvipath.com) before you start.

IMPORTANT SAFETY INFORMATION AND USE

What should I tell my healthcare provider before taking YORVIPATH? (cont'd)

Before using YORVIPATH, tell your healthcare provider about all of your medical conditions, including if you:

- are pregnant or plan to become pregnant. It is not known if YORVIPATH will harm your unborn baby. Tell your healthcare provider if you become pregnant during treatment with YORVIPATH.
- are breastfeeding or plan to breastfeed. It is not known if YORVIPATH passes into your breast milk. Talk to your healthcare provider about the best way to feed your baby during treatment with YORVIPATH.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. YORVIPATH and other medicines may affect each other causing side effects.

Please see Important Safety Information throughout. Please click to read or print the [Prescribing Information](#) and [Medication Guide](#) for YORVIPATH.

Yorvipath®
palopegteriparatide
Injection 300 mcg/mL

Get personalized support at each step in your treatment journey



Once enrolled in A-S-A-P, you will have access to a team of caring professionals providing you with support throughout your treatment journey, including:



Understanding insurance coverage



Help seeking financial assistance to pay for YORVIPATH



Refills and ongoing support to stay on track with YORVIPATH



Connecting you to a Patient Access Liaison (PAL)

Have any questions? We are here to help!

To talk to a member of our support team or get connected to a PAL, call [1-844-442-7236](tel:1-844-442-7236) (8 AM to 8 PM ET, Monday through Friday).

IMPORTANT SAFETY INFORMATION AND USE

What are the possible side effects of YORVIPATH?

YORVIPATH may cause serious side effects:

- **Decrease in blood pressure when you change positions (orthostatic hypotension).** Some people may feel dizzy, get a fast heartbeat, or feel light-headed right after injecting YORVIPATH. For the first few doses, give your injection of YORVIPATH in a place where you can sit or lie down right away if you get these symptoms. If your symptoms get worse or do not go away, contact your healthcare provider before you continue using YORVIPATH.

Please see Important Safety Information throughout. Please click to read or print the [Prescribing Information](#) and [Medication Guide](#) for YORVIPATH.

Yorvipath[®]
palopegteriparatide
Injection 300 mcg/mL



Your PAL can help you with many things, such as learning about the insurance process, how YORVIPATH fits into your treatment, and staying on track with your treatment journey.



“It is important to take things one step at a time and find a strong community that understands.”

– Denise, **actual patient**



Hear more stories from people like you at [YORVIPATH.com](https://www.yorvipath.com).

IMPORTANT SAFETY INFORMATION AND USE

What are the possible side effects of YORVIPATH?

YORVIPATH may cause serious side effects: (cont'd)

- Digoxin toxicity if you are using digoxin and YORVIPATH at the same time. Tell your healthcare provider if you experience irregular heart rhythm, heart palpitations, confusion, loss of appetite, nausea, vomiting, diarrhea, or vision problems.

Please see Important Safety Information throughout. Please click to read or print the [Prescribing Information](#) and [Medication Guide](#) for YORVIPATH.

Yorvipath[®]
palopegteriparatide
Injection 300 mcg/mL

Take charge with YORVIPATH

With YORVIPATH, you can expect:



24-hr delivery of PTH



A once-daily injection



Individualized dosing*



Ongoing support from A·S·A·P



Learn more about
taking YORVIPATH
at [YORVIPATH.com](https://www.yorvipath.com)

*Your doctor will support you through the titration process to find your individualized dose.
Actual patients were compensated by Ascendis Pharma.

IMPORTANT SAFETY INFORMATION AND USE

What are the possible side effects of YORVIPATH? (cont'd)

The most common side effects of YORVIPATH include injection site reactions, vasodilatory signs or symptoms such as a drop in blood pressure or lightheadedness when standing up, headache, diarrhea, back pain, high calcium levels, and sore throat.

These are not all of the possible side effects of YORVIPATH. Call your doctor for medical advice about side effects. You are encouraged to report side effects to FDA at [1-800-FDA-1088](https://www.fda.gov/medwatch) or at www.fda.gov/medwatch. You may also report side effects to Ascendis Pharma at [1-844-442-7236](https://www.ascendispharma.com).

Please see Important Safety Information throughout. Please click to read or print the [Prescribing Information](#) and [Medication Guide](#) for YORVIPATH.

Yorvipath[®]
palopegteriparatide
Injection 300 mcg/mL



© August 2026 Ascendis Pharma Endocrinology, Inc. All rights reserved.
YORVIPATH[®], Ascendis Signature Access Program[®], Ascendis[®], the
Ascendis Pharma logo and the company logo are trademarks owned
by the Ascendis Pharma Group. US-COMMPH-2600210 08/26